

SELF DELTA-EQUIVALENCE FOR LINKS WHOSE MILNOR'S ISOTOPY INVARIANTS VANISH

AKIRA YASUHARA

ABSTRACT. For an n -component link L , the Milnor's isotopy invariant is defined for each multi-index $I = i_1 i_2 \dots i_m$ ($i_j \in \{1, \dots, n\}$). Here m is called the length. Let $r(I)$ denote the maximum number of times that any index appears. It is known that Milnor invariants with $r = 1$ are link-homotopy invariant. N. Habegger and X. S. Lin showed that two string links are a link-homotopy if and only if their Milnor invariants with $r = 1$ coincide. This gives us that a link in S^3 is link-homotopic to a trivial link if and only if the all Milnor invariants of the link with $r = 1$ vanish. Although Milnor invariants with $r = 2$ are not link-homotopy invariants, T. Fleming and the author showed that Milnor invariants with $r \leq 2$ are self Δ -equivalence invariants. In this paper, we give a self Δ -equivalence classification of the set of n -component links in S^3 whose Milnor invariants with length $\leq 2n - 1$ and $r \leq 2$ vanish. As a corollary, we have that a link is self Δ -equivalent to a trivial link if and only if the all Milnor invariants of the link with $r \leq 2$ vanish. This is a geometric characterization for links whose Milnor invariants with ≤ 2 vanish. The chief ingredient in our proof is Habiro's clasper theory. We also give an alternate proof of a link-homotopy classification of string links by using clasper theory.

1. INTRODUCTION

For an n -component link L , *Milnor invariant* $\overline{\mu}_L(I)$ is defined for each multi-index $I = i_1 i_2 \dots i_m$ ($i_j \in \{1, \dots, n\}$) [17, 18]. Here m is called the *length* of $\overline{\mu}_L(I)$ and denoted by $|I|$. Let $r(I)$ denote the maximum number of times that any index appears. For example, $r(1123) = 2$, $r(1231223) = 3$. It is known that if $r(I) = 1$, then $\overline{\mu}_L(I)$ is a *link-homotopy* invariant [17], where link-homotopy is an equivalence relation on links generated by self crossing changes. Similarly, for a string link L , Milnor invariant $\mu_L(I)$ is defined [8]. While Milnor invariants are not strong enough to give a link-homotopy classification for links, they are complete for string links. In fact, the following is known [8].

Theorem 1.1 ([8]). *Two n -component string links L and L' are link-homotopic if and only if $\mu_L(I) = \mu_{L'}(I)$ for any I with $r(I) = 1$.*

We will give an alternate proof in section 4 via clasper theory. Actually we will give representatives determined by Milnor link-homotopy invariants for the link-homotopy classes explicitly, see Theorem 4.3. As a corollary, we have that for n -component string links L and L' , and for a positive integer k ($k \leq n$), $\mu_L(I) = \mu_{L'}(I)$ for any I

2000 *Mathematics Subject Classification.* 57M25, 57M27.

Key words and phrases. Δ -move, self Δ -move, C_n -move, link-homotopy, self Δ -equivalence, Milnor invariant, string link, Brunnian link, clasper.

The author is partially supported by a Grant-in-Aid for Scientific Research (C) (#18540071) of the Japan Society for the Promotion of Science.

with $r(I) = 1$ and $|I| \leq k$ if and only if L and L' are transformed into each other by combining link-homotopies and C_k -moves, see Corollary 4.5.

For a string link L , let $\text{cl}(L)$ denote the *closure* of L . It follows from the definitions that $\mu_L(I) = \overline{\mu}_{\text{cl}(L)}(I)$ if $\mu_L(J) = 0$ for any J with $|J| < |I|$. Since the Milnor invariants of trivial (string) links are 0, this and Theorem 1.1 imply the following. The proposition below also follows from Milnor's link-homotopy classification theorem for Brunnian links [17].

Proposition 1.2 ([17, Section 5]). *A link L in S^3 is link-homotopic to a trivial link if and only if $\overline{\mu}_L(I) = 0$ for any I with $r(I) = 1$.*

Although Milnor invariants with $r \geq 2$ are not necessarily link-homotopy invariants, they are generalized link-homotopy invariants. In fact, Fleming and the author [3] showed that Milnor invariants with $r \leq k$ are *self C_k -equivalence* invariants, where the (self) C_k -equivalence is an equivalence relation on (string) links generated by (self) C_k -moves defined as follows.

A C_n -move is a local move on (string) links as illustrated in Figure 1.1. (A C_1 -move is defined as the crossing change). These local moves were introduced by Habiro [9]. A C_n -move is called a *self C_n -move* if the all strands in Figure 1.1 belong to the same component of a (string) link [25].




FIGURE 1.1. A C_n -move involves $n + 1$ strands of a link.

The C_n -move (resp. self C_n -move) generates an equivalence relation on links, called the C_n -equivalence (resp. self C_n -equivalence). This notion can also be defined by using the theory of claspers (see section 2). The (self) C_n -equivalence relation becomes finer as n increases, i.e., the (self) C_m -equivalence implies the (self) C_k -equivalence for $m > k$. We remark that (self) C_2 -move is same as (self) Δ -move defined by [20]. The Δ -move is defined as a local move as illustrated in Figure 1.2. We call the (self) C_2 -equivalence the (self) Δ -equivalence.

A self Δ -equivalence classification of 2-component links was shown by Y. Nakanishi and Y. Ohyama [21]. It is still open for links with at least 3 components. Here we give the following theorem.




FIGURE 1.2.

Theorem 1.3. *Let L and L' be n -component links. Suppose that $\overline{\mu}_L(I) = \overline{\mu}_{L'}(I) = 0$ for any I with $|I| \leq 2n - 1$ and $r(I) \leq 2$. Then L and L' are self Δ -equivalent if and only if $\overline{\mu}_L(J) = \overline{\mu}_{L'}(J)$ for any J with $|J| = 2n$ and $r(J) = 2$.*

Remark 1.4. (1) The ‘only if’ part follows directly from the fact that Milnor invariants with $r \leq k$ are self C_k -equivalence invariants [3].

(2) In the last section, we characterize n -component links whose Milnor invariants of length $\leq 2n - 1$ and $r \leq 2$ vanish. More precisely, the Milnor invariants of an n -component link with length $\leq 2n - 1$ and $r \leq 2$ vanish if and only if, for any integer i in $\{1, \dots, n\}$, it is self Δ -equivalent to a Brunnian link L_i such that the i th component K of L_i is null-homotopic in $S^3 \setminus (L_i - K)$ (Theorem 6.3). As an example, we will give a 3-component Brunnian link $L = K_1 \cup K_2 \cup K_3$ such that K_1 is not null-homotopic in $S^3 \setminus (L - K_1)$ and K_i is null-homotopic in $S^3 \setminus (L - K_i)$ ($i = 2, 3$) (Example 6.4). In particular, L is link-homotopic to a trivial link. There is no such a link with 2 components, i.e., if a 2-component link is link-homotopic to a trivial link, then it is self Δ -equivalent to a Brunnian link $K_1 \cup K_2$ such that K_i is null-homotopic in $S^3 \setminus K_j$ ($\{i, j\} = \{1, 2\}$).

For 2-component links, Proposition 1.2 can be generalized [21]. Theorem 1.3 gives us the following corollary which is a generalization of Proposition 1.2 for links with arbitrarily many components. This gives us a geometric characterization for links whose Milnor invariants with ≤ 2 vanish.

Corollary 1.5. *A link L is self Δ -equivalent to a trivial link if and only if $\overline{\mu}_L(I) = 0$ for any I with $r(I) \leq 2$.*

Remark 1.6. (1) This corollary gives an affirmative answer for an open question remained in [3].

(2) For string links, Corollary 1.5 does not hold, i.e., there are 2-string links such that their Milnor invariants $\mu(I)$ with $r(I) \leq 2$ vanish and they are not self Δ -equivalent to a trivial string link [4].

(3) Since C_k -move ($k \geq 3$) is not unknotting operation, it is impossible to generalize the corollary above. It is reasonable to consider the following question: If $\overline{\mu}_L(I) = 0$ for any I with $r(I) \leq k$, then is L self C_k -equivalent to a completely split link? Fleming and the author gave a negative answer to the question [3]. In fact, there is a 2-component boundary link L such that L is not self C_3 -equivalent to a split link. Note that the all Milnor invariants of a boundary link vanish.

By combining Lemma 3.2 ([18, Theorem 7]), Proposition 1.2 and Corollary 1.5, we have the following corollary.

Corollary 1.7. *Let L be an n -component link and let $L(2)$ be a $2n$ -component link obtained from L by replacing each component of L with zero framed 2 parallel copies of it. Then L is self Δ -equivalent to a trivial link if and only if $L(2)$ is link-homotopic to a trivial link.*

Remark 1.8. For an n -component link, let $L(k)$ be a kn -component link obtained from L by replacing each component of L with zero framed k parallel copies of it. In the proof of Theorem 2.1 in [3], it is shown that if two links L and L' are self

C_k -equivalent, then $L(k)$ and $L'(k)$ are link homotopic. So one might expect that if $L(2)$ and $L'(2)$ are link homotopic, then L and L' are self Δ -equivalent. But this is not true. The reason is the following: There are 2-component links L and L' such that they are concordant and are not self Δ -equivalent [22], [23]. The fact that L and L' are concordant implies that $L(2)$ and $L'(2)$ are concordant. Since link-concordance implies link-homotopy [5], [6], $L(2)$ and $L'(2)$ are link-homotopic.

An n -component link $L = K_1 \cup \cdots \cup K_n$ is called a *boundary link* if there is a disjoint union $X = F_1 \cup \cdots \cup F_n$ of orientable surfaces such that $\partial X = L$ and $\partial F_i = K_i$ ($i = 1, 2, \dots, n$). An n -component link L is called a *homology boundary link* if $\pi_1(S^3 \setminus L)$ admits an epimorphism from $\pi_1(S^3 \setminus L)$ to a free group of rank n [27]. An every boundary link is a homology boundary link. T. Shibuya and the author showed that all boundary links are self Δ -equivalent to trivial links [26]. In [24], Shibuya showed that all ribbon links are self Δ -equivalent to trivial links.

Whether the homology boundary links are self Δ -equivalent to trivial links and whether the slice links are self Δ -equivalent to trivial links have remained as open questions. Since all Milnor invariants of homology boundary links vanish, and since Milnor invariants are concordance invariants, we have the following corollary, which give affirmative answers for the open questions.

Corollary 1.9. *If L is concordant to a homology boundary link, then L is self Δ -equivalent to a trivial link.*

Acknowledgments. The author would like to thank Jean-Baptiste Meilhan for many useful discussions. The first joint work [16] leads to this work. He is also very grateful to Professor Tim Cochran for helpful comments.

2. CLASPER

Let us briefly recall from [10] the basic notions of clasper theory for (string) links. In this paper, we essentially only need the notion of C_k -tree. For a general definition of clasps, we refer the reader to [10].

Definition 1. Let L be a link in S^3 (resp. a string link in $D^2 \times I$). An embedded disk F in S^3 (resp. $D^2 \times I$) is called a *tree clasper* for L if it satisfies the following (1), (2) and (3):

- (1) F is decomposed into disks and bands, called *edges*, each of which connects two distinct disks.
- (2) The disks have either 1 or 3 incident edges, called *leaves* or *nodes* respectively.
- (3) L intersects F transversely and the intersections are contained in the union of the interior of the leaves.

The *degree* of a tree clasper is the number of the leaves *minus* 1. (In [10], a tree clasper and a leaf are called a *strict tree clasper* and a *disk-leaf* respectively.) A degree k tree clasper is called a C_k -tree (or a C_k -clasper). A C_k -tree is *simple* if each leaf intersects L at one point.

We will make use of the drawing convention for clasps of [10, Fig. 7], except for the following: $\oplus$ (resp. $\ominus$) on an edge represents a positive (resp. negative) half-twist. (This replaces the convention of a circled S (resp. S^{-1}) used in [10]).

Given a C_k -tree T for a link L in S^3 , there is a procedure to construct a framed link $\gamma(T)$ in a regular neighborhood of T . *Surgery along T* means surgery along $\gamma(T)$. Since there exists a canonical homeomorphism between S^3 and the manifold $S^3_{\gamma(T)}$, surgery along the C_k -tree T can be regarded as a local move on L in S^3 . We say that the resulting link L_T in S^3 is *obtained by surgery along T* . In particular, surgery along a simple C_k -tree illustrated in Figure 2.1 is equivalent to band-summing a copy of the $(k+1)$ -component Milnor link (see [17, Fig. 7]), and is equivalent to a C_k -move as defined in the introduction (Figure 1.1). Similarly, for a disjoint union of trees $T_1 \cup \dots \cup T_m$, we can define $L_{T_1 \cup \dots \cup T_m}$ as a link by surgery along $T_1 \cup \dots \cup T_m$. A C_k -tree T having the shape of the tree clasper in Figure 2.1 is called *linear*, and the left-most and right-most leaves of T in Figure 2.1 are called *ends* of T . Ends of T are not uniquely determined. There are 4 choices for an each linear tree.




FIGURE 2.1. Surgery along a simple C_k -tree.

It is known that the C_k -equivalence as defined in section 1 coincides with the equivalence relation on links generated by surgery along C_k -trees and ambient isotopies. Two (string) links L and L' are C_k -equivalent if and only if there is a disjoint union of simple C_k -trees $G_1 \cup \dots \cup G_m$ such that L' is ambient isotopic to $L_{G_1 \cup \dots \cup G_m}$ [10, Theorem 3.17].

Definition 2. Let $L = K_1 \cup \dots \cup K_n$ be an n -component (string) link. A (simple) C_k -tree T for L is a (simple) C_k^a -tree (resp. C_k^d -tree, C_k^s -tree) if it satisfies the following:

- (1) For each disk-leaf f of T , $f \cap L$ is contained in a single component of L , and
- (2) $|\{i \mid T \cap K_i \neq \emptyset\}| = n$ (resp. $= k+1, 1$).

Note that n is the number of the components of L and that $k+1$ is the number of leaves of T . If T is simple, T always satisfies the condition (1). The C_k^* -equivalence ($*$ = a, d, s) is an equivalence relation on (string) links generated by surgery along C_k^* -trees and ambient isotopies. Note that C_k^s -equivalence is same as self C_k -equivalence. For a simple C_k -tree T , the set $\{i \mid T \cap K_i \neq \emptyset\}$ is called *index* of T , and denote it by $\text{index}(T)$. And let $r_i(T)$ be the number of intersection points in $T \cap K_i$ ($i = 1, \dots, n$). The $(C_l^s + C_k^*)$ -equivalence ($C_k^* = C_k, C_k^a$, or C_k^d) is an equivalence relation on (string) links generated by surgery along C_l^s - or C_k^* -trees. By the arguments similar to that in the proof of [10, Theorem 3.17], we have that two (string) links L and L' are $(C_l^s + C_k^*)$ -equivalent if and only if there is a disjoint union of simple C_l^s - or C_k^* -trees $T_1 \cup \dots \cup T_m$ such that L' is ambient isotopic to $L_{T_1 \cup \dots \cup T_m}$. We use the notation

$L \stackrel{C_k^*}{\sim} L'$ (resp. $L \stackrel{C_l^s + C_k^*}{\sim} L'$) for C_k^* -equivalent (resp. $(C_l^s + C_k^*)$ -equivalent) links L and L' .

Recall that a string link is a tangle without closed components (see [8] for a precise definition). The set of ambient isotopy classes of the n -component string links has a monoid structure with composition given by the *stacking product*, denoted by $*$, and with the trivial n -component string link $\mathbf{1}_n$ as unit element.

In the following, we give several lemmas. The proofs of Lemmas 2.1, 2.2 and 2.3 are essentially given in [10] (see also section 1.4 in [14]), and Lemma 2.4 essentially shown in [7] (see also [2], [15]), while they did not care about r_j of claspers in [2], [7], [10], [14], [15]. If we follow their proofs with paying attention to r_j , we will see the proof of Lemmas 2.1, 2.2, 2.3 and 2.4.

Lemma 2.1 (cf. [10, Propositions 4.5, 4.6]). *Let T be a simple C_k -tree for an n -component (string) link L , and let T' (resp. T'' , and T''') be obtained from T by changing a crossing of an edge and the i th component K_i of L (resp. an edge of T , and an edge of another simple clasper G) (see Figure 2.2). Then*

- (1) $L_T \stackrel{C_{k+1}}{\sim} L_{T'}$, and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with indices $\text{index}(T) \cup \{i\}$ and $r_j \geq r_j(T)$ ($j = 1, \dots, n$).
- (2) $L_T \stackrel{C_{k+1}}{\sim} L_{T''}$, $L_{T \cup G} \stackrel{C_{k+1}}{\sim} L_{T'' \cup G}$, and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with $r_j \geq r_j(T)$ ($j = 1, \dots, n$).




FIGURE 2.2.

Lemma 2.2 (cf. [10, Propositions 4.4]). *Let T_1 (resp. T_2) be a simple C_k -tree (resp. C_l -tree) for an n -component (string) link L , and let T'_1 be obtained from T_1 by sliding a leaf of T_1 over a leaf of T_2 (see Figure 2.3). Suppose that $k \geq l$. Then $L_{T_1 \cup T_2} \stackrel{C_{k+1}}{\sim} L_{T'_1 \cup T_2}$, and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with $r_j \geq r_j(T_1)$ ($j = 1, \dots, n$).*




FIGURE 2.3. Sliding a leaf over another leaf.

Lemma 2.3 (cf. [10, Claim in p-36]). *Let T be a simple C_k -tree for $\mathbf{1}_n$ and let $\bar{T}$ be a simple C_k -trees obtained from T by adding a half-twist on an edge. Then $(\mathbf{1}_n)_T * (\mathbf{1}_n)_{\bar{T}} \stackrel{C_{k+1}}{\sim} \mathbf{1}_n$, and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with $r_j \geq r_j(T)$ ($j = 1, \dots, n$).*

Lemma 2.4 (cf. [7, Theorem 6.7], [15, Lemma 2.9]). *Consider simple C_k -trees T_I , T_H and T_X for $\mathbf{1}_n$ which differ only in a small ball as illustrated in Figure 2.4. Then $(\mathbf{1}_n)_{T_I} \stackrel{C_{k+1}}{\sim} (\mathbf{1}_n)_{T_H} * (\mathbf{1}_n)_{T_X}$, and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with $r_j \geq r_j(T_I)$ ($j = 1, \dots, n$).*




FIGURE 2.4. The IHX relation for C_k -trees.

By combining the proof of [10, Claim in p-26] and [10, Propositions 4.4, 4.5 and 4.6], we have the following.

Lemma 2.5 (cf. [10, Claim in p-26]). *Let G be a C_k -tree for $\mathbf{1}_n$. Let f_1 and f_2 be two disks obtained by splitting a leaf f of G along an arc α as shown in Figure 2.5 (i.e., $f = f_1 \cup f_2$ and $f_1 \cap f_2 = \alpha$). Then, $(\mathbf{1}_n)_G \stackrel{C_{k+1}}{\sim} (\mathbf{1}_n)_{G_1} * (\mathbf{1}_n)_{G_2}$, where G_i denotes the C_k -tree for $\mathbf{1}_n$ obtained from G by replacing f with f_i ($i = 1, 2$).*




FIGURE 2.5. Splitting a leaf.

An n -component (string) link L is *Brunnian* if every proper sublink of L is trivial. In particular, any trivial (string) link is Brunnian. The n -component Brunnian (string) links are characterized by C_{n-1}^a -equivalence as follows.

Proposition 2.6 ([11, 19]). *Let L be an n -component (string) link in S^3 . Then L is Brunnian if and only if L is obtained from a trivial (string) link by surgery along simple C_{n-1}^a -trees.*

By the argument similar to that in the proof of Theorem 1.2 in [19], we have the following lemma. In [19], they proved it with using ‘band description’ defined by K. Taniyama and the author [28]. Here we give a proof with using clasper.

Lemma 2.7 (cf. [19, Theorem 1.2]). *Let L be an n -component Brunnian link in S^3 . If L is obtained from a trivial link O by surgery along C_1^s -trees with indices $\{i\}$, then L is obtained from O by surgery along simple C_n^a -trees with $r_i = 2$.*

Proof. Set $O = O_1 \cup \cdots \cup O_n$. It is enough to consider the case when $i = 1$. There is a disjoint union F_1 of simple C_1^s -trees with indices $\{1\}$ such that $L = O_{F_1}$. Note that $r_1 = 2$ for all C_1^s -trees in F_1 .

Since L is Brunnian, $L \setminus O_2$ is trivial. This implies that a split sum of $L \setminus O_2$ and O_2 is trivial. Hence L can be deformed into trivial by crossing changes between O_2 and edges of C_1^s -trees of F_1 . By Lemma 2.1, we have that L is obtained from O by surgery along a disjoint union F_2 of simple C_2 -trees with indices $\{1, 2\}$ and $r_1 = 2$. So we have $L = O_{F_2}$.

Since $L \setminus O_3$ is trivial, L can be deformed into trivial by crossing changes between O_3 and edges of C_2 -trees in F_2 . By Lemma 2.1, there is a disjoint union F_3 of simple C_3 -trees with indecis $\{1, 2, 3\}$ and $r_1 = 2$ such that $L = O_{F_3}$.

Repeating this step, we have that there is a disjoint union F_n of simple C_n -trees with indices $\{1, \dots, n\}$ and $r_1 = 2$ such that $L = O_{F_n}$. This completes the proof. $\square$

By the arguments similar to that in the proof of [3, Proposition 3.1], we have

Proposition 2.8 (cf. [3, Proposition 3.1]). *A (string) link L' is obtained from L by surgery along a simple C_l -tree with leaves $f_1, f_2, \dots, f_{l+1}$, then for any k ($1 \leq k < l$) and any subset $\{w_1, \dots, w_{k+1}\} \subset \{1, \dots, l+1\}$, there are simple C_k -trees T_j ($j = 1, \dots, m$) with leaves $f_{j1}, \dots, f_{j(k+1)}$ such that f_{ji} and f_{wi} grasp the same component of L for each $i (= 1, \dots, k+1)$, and that L' is obtained from L by surgery along $T_1, \dots, T_m$.*

A simple C_k -tree T is a $C_k^{(l)}$ -tree if $\max\{r_j(T) \mid j = 1, \dots, n\} \geq l$. Two links L and L' are $C_k^{(l)}$ -equivalent if L is obtained from L' by ambient isotopy and surgery along simple $C_k^{(l)}$ -trees. The following proposition is a corollary of Proposition 2.8.

Proposition 2.9 (cf. [3, Proposition 3.1]). *If two (string) links are $C_n^{(k+1)}$ -equivalent, then they are self C_k -equivalent. Moreover, for some i , if the $C_n^{(k+1)}$ -equivalence is realized by surgery along simple $C_n^{(k+1)}$ -trees with $r_i \geq k+1$, then the self C_k -equivalence is realized by surgery along simple C_k^s -trees with indices $\{i\}$.*

3. MILNOR INVARIANTS

J. Milnor defined in [17] a family of invariants of oriented, ordered links in S^3 , known as Milnor's $\bar{\mu}$ -invariants.

Given an n -component link L in S^3 , denote by G the fundamental group of $S^3 \setminus L$, and by G_q the q th subgroup of the lower central series of G . We have a presentation of G/G_q with n generators, given by a meridian m_i of the i th component of L . So for $1 \leq i \leq n$, the longitude l_i of the i th component of L is expressed modulo G_q as a word in the m_i 's (abusing notations, we still denote this word by l_i).

The Magnus expansion $E(l_i)$ of l_i is the formal power series in non-commuting variables $X_1, \dots, X_n$ obtained by substituting $1 + X_j$ for m_j and $1 - X_j + X_j^2 - X_j^3 + \cdots$ for m_j^{-1} , $1 \leq j \leq n$.

Let $I = i_1 i_2 \dots i_{k-1} j$ ($k \leq q$) be a multi-index (i.e., a sequence of possibly repeating indices) among $\{1, \dots, n\}$. Denote by $\mu_L(I)$ the coefficient of $X_{i_1} \cdots X_{i_{k-1}}$ in the Magnus expansion $E(l_j)$. Milnor invariant $\bar{\mu}_L(I)$ is the residue class of $\mu_L(I)$ modulo

the greatest common divisor of all Milnor invariants $\mu_L(J)$ such that J is obtained from I by removing at least one index. As we mentioned in section 1, $|I| = k$ is called the length of Milnor invariant $\overline{\mu}_L(I)$.

The indeterminacy comes from the choice of the meridians m_i . Equivalently, it comes from the indeterminacy of representing the link as the closure of a string link [8]. Indeed, $\mu(I)$ is a well-defined invariant for string links.

The following 4 lemmas play an important roles in calculating Milnor invariants.

Lemma 3.1 ([17, section 5]). *Let $M_n = K_1 \cup \dots \cup K_n$ be the n -component Milnor link as illustrated in Figure 3.1. Then the Milnor invariants of length $\leq n-1$ vanish, and*

$$\overline{\mu}_{M_n}(i_1 i_2 \dots i_{n-2} \ n-1 \ n) = \begin{cases} 1 & \text{if } i_1 i_2 \dots i_{n-2} = 12 \dots n-2, \\ 0 & \text{otherwise.} \end{cases}$$




FIGURE 3.1. Milnor link

Lemma 3.2 ([18, Theorem 7]). *Let L' be a link obtained from a link L by taking the appropriate number of zero framed parallels of the components of L . Suppose the i th component of L' corresponds to the $h(i)$ th component of L , then*



$$\overline{\mu}_{L'}(i_1 i_2 \dots i_m) = \overline{\mu}_L(h(i_1) h(i_2) \dots h(i_m)).$$

Lemma 3.3 ([16, Lemma 3.3]). *Let L and L' be n -component string links such that all Milnor invariants of L (resp. L') of length $\leq m$ (resp. $\leq m'$) vanish. Then $\mu_{L*L'}(I) = \mu_L(I) + \mu_{L'}(I)$ for all I of length $\leq m + m'$.*

Lemma 3.4 ([10, Theorem 7.2]). *The Milnor invariants of length $\leq k$ for (string) links are invariants of the C_k -equivalence.*

4. LINK-HOMOTOPY OF STRING LINKS

Let $\pi : \{1, \dots, k\} \longrightarrow \{1, \dots, n\}$ ($k \leq n$) be an injection such that $\pi(i) < \pi(k-1) < \pi(k)$ ($i \in \{1, \dots, k-2\}$), and let $\mathcal{F}_k$ be the set of such injections. For $\pi \in \mathcal{F}_k$, let T_π and $\overline{T}_\pi$ be simple C_{k-1}^d -trees as illustrated in Figure 4.1, and set $V_\pi = (\mathbf{1}_n)_{T_\pi}$ and $V_\pi^{-1} = (\mathbf{1}_n)_{\overline{T}_\pi}$. Here, Figure 4.1 are the images of homeomorphisms from the neighborhoods of T_π and $\overline{T}_\pi$ to the 3-ball. Although V_π and V_π^{-1} are not unique up to ambient isotopy, by Lemmas 2.1 and 2.3, it is unique up to C_k -equivalence. So, for any $\pi \in \mathcal{F}_k$, we may choose V_π and V_π^{-1} uniquely up to C_k -equivalence. In

FIGURE 4.1. $\pi(i)$ means $\pi(i)$ th component of $\mathbf{1}_n$ FIGURE 4.2. Milnor link $M_\pi = K_{\pi(1)} \cup \cdots \cup K_{\pi(k)}$

particular, we may choose V_π so that $\text{cl}(V_\pi)$ is the Milnor link $M_\pi = K_{\pi(1)} \cup \cdots \cup K_{\pi(k)}$ as illustrated in Figure 4.2 (cf. Figure 2.1).

For $\pi \in \mathcal{F}_k$, set

$$\mu_\pi(L) = \mu_L(\pi(1)\pi(2)\cdots\pi(k))$$

By Lemma 3.1, we have the following lemma.

Lemma 4.1. *For any $\pi, \pi' \in \mathcal{F}_k$,*

$$\mu_\pi(V_{\pi'}) = \begin{cases} 1 & \text{if } \pi = \pi', \\ 0 & \text{if } \pi \neq \pi', \end{cases}$$

and the Milnor invariants of $V_{\pi'}$ of length $\leq k-1$ vanish.

Lemma 4.2. *Let T be a simple C_{k-1}^d -tree (resp. C_{n-1}^a -tree) for an n -component string link L . Then L_T is C_k -equivalent (resp. C_n^a -equivalent) to $L * L'$, where*

$$L' = \prod_{\pi \in \mathcal{F}_k} V_\pi^{\mu_\pi(L_T) - \mu_\pi(L)}.$$

Proof. Suppose that T is a simple C_{k-1}^d -tree. By Lemma 2.1, L_T is C_k -equivalent to $L * (\mathbf{1}_n)_{T'}$, where T' is a simple C_{k-1}^d -tree. Set $\text{index}(T') = \{i_1, \dots, i_k\}$ ($i_j < i_{j+1}$). Consider induction on the length of the path connecting the two leaves grasping i_{k-1} th and i_k th components of $\mathbf{1}_n$, and apply Lemma 2.4, we have that $(\mathbf{1}_n)_{T'}$ is C_k -equivalent to a string link which is obtained from $\mathbf{1}_n$ by surgery along simple C_{k-1}^d -trees whose ends grasp i_{k-1} th and i_k th components of $\mathbf{1}_n$. By Lemmas 2.1, 2.2 and 2.3, we have that

$$(\mathbf{1}_n)_{T'} \stackrel{C_k}{\sim} L'' = \prod_{\pi \in \mathcal{F}_k} V_\pi^{x_\pi}.$$

By Lemmas 3.4, 3.3 and 4.1,

$$\begin{aligned} \mu_{\pi'}(L_T) = \mu_{\pi'}(L * L'') &= \mu_{\pi'}(L) + \mu_{\pi'}(L'') \\ &= \mu_{\pi'}(L) + \sum_{\pi \in \mathcal{F}_k} x_\pi \mu_{\pi'}(V_\pi) = \mu_{\pi'}(L) + x_{\pi'} \cdot \end{aligned}$$

If T is a simple C_{n-1}^a -tree, the arguments similar to that in the above can be applied. And we have the conclusion. $\square$

The following theorem gives representatives, which depend on only Milnor invariants, for the link-homotopy classes.

Theorem 4.3. *Let L be an n -component string link. Then L is link-homotopic to $L_1 * L_2 * \cdots * L_{n-1}$, where*

$$L_i = \prod_{\pi \in \mathcal{F}_{i+1}} V_{\pi}^{x_{\pi}}, \quad x_{\pi} = \begin{cases} \mu_L(\pi(1)\pi(2)) & \text{if } i = 1, \\ \mu_L(\pi(1)\dots\pi(i+1)) - \mu_{L_1\dots L_{i-1}}(\pi(1)\dots\pi(i+1)) \\ \quad (= \mu_{L_i}(\pi(1)\dots\pi(i+1))) & \text{if } i \geq 2. \end{cases}$$

Remark 4.4. The presentation $L_1 * L_2 * \cdots * L_{n-1}$ of L depends on the choice of order on the elements in $\mathcal{F}_i$ ($i = 2, \dots, n$). If we put $\mathcal{F}_2 \cup \cdots \cup \mathcal{F}_n = \{\pi_1, \dots, \pi_q\}$ so that for $i < j$, any element in $\mathcal{F}_i$ appears before the elements in $\mathcal{F}_j$, then by Theorem 4.3 and Lemmas 3.3 and 4.1, L is link-homotopic to $V_{\pi_1}^{x_{\pi_1}} * \cdots * V_{\pi_q}^{x_{\pi_q}}$ ($x_{\pi_k} = \mu_{\pi_k}(L) - \mu_{\pi_k}(\prod_{i=1}^{k-1} V_{\pi_i}^{x_{\pi_i}})$). Note that the representation is unique up to link-homotopy.

Proof. Since C_1 -move is the crossing change, L is C_1 -equivalent to the trivial string link $\mathbf{1}_n$. So L is obtained from $\mathbf{1}_n$ by surgery along simple C_1 -trees.

Note that a simple C_1 -tree is either a simple C_1^s -tree or a simple C_1^d -tree, and that C_1^s -equivalence preserves the value of $\mu(I)$ for any I with $r(I) = 1$. Since L is C_1^s -equivalent to a link which is obtained from $\mathbf{1}_n$ by surgery along C_1^d -trees, by Lemmas 4.2, 2.1, 2.2 and 2.3,

$$L \stackrel{C_1^s + C_2}{\sim} \prod_{\pi \in \mathcal{F}_2} V_{\pi}^{\mu_{\pi}(L)} (= L_1).$$

A C_k -tree ($k \geq 2$) is either a $C_k^{(2)}$ -tree or a C_k^d -tree, and a $C_k^{(2)}$ -equivalence implies C_1^s -equivalence (Proposition 2.9), and hence $(C_1^s + C_k)$ -equivalence implies $(C_1^s + C_k^d)$ -equivalence. So L is obtained from L_1 by surgery along simple C_1^s - and C_2^d -trees.

By Lemmas 4.2, 2.1, 2.2 and 2.3,

$$L \stackrel{C_1^s + C_3}{\sim} L_1 * \prod_{\pi \in \mathcal{F}_3} V_{\pi}^{\mu_{\pi}(L) - \mu_{\pi}(L_1)} (= L_1 * L_2).$$

Therefore L and $L_1 * L_2$ are $(C_1^s + C_3^d)$ -equivalent.

Repeating these processes, we have that

$$L \stackrel{C_1^s + C_n}{\sim} L_1 * L_2 * \cdots * L_{n-1}.$$

Since any simple C_n -tree for an n -component string link is a $C_n^{(2)}$ -tree, $(C_1^s + C_n)$ -equivalence implies C_1^s -equivalence, i.e., link-homotopy. $\square$

By Theorem 4.3, we have the following corollary.

Corollary 4.5. *For a natural number $k(\leq n)$, n -component string links L and L' are $(C_1^s + C_k)$ -equivalent if and only if $\mu_L(I) = \mu_{L'}(I)$ for any I with $r(I) = 1$ and $|I| \leq k$.*

Proof. The ‘only if’ part follows from Lemma 3.4. Now we will prove ‘if’ part.

By Theorem 4.3, L and L' are link-homotopic to $L_1 * L_2 * \cdots * L_{n-1}$ and $L'_1 * L'_2 * \cdots * L'_{n-1}$ respectively. Note that both L_i and L'_i are C_i -equivalent to $\mathbf{1}_n$. So L and L' are $(C_1^s + C_k)$ -equivalent to $L_1 * L_2 * \cdots * L_{k-1}$ and $L'_1 * L'_2 * \cdots * L'_{k-1}$ respectively. Since $\mu_L(I) = \mu_{L'}(I)$ for any I with $r(I) = 1$ and $|I| \leq k$, $L_i = L'_i$ ($i = 1, \dots, k-1$). This completes the proof. $\square$

Theorem 1.1 follows directly from Corollary 4.5.

Proof of Theorem 1.1. It is enough to show ‘if’ part. Since a C_n -move for an n -component string link is a $C_n^{(2)}$ -move, by Proposition 2.9, C_n -equivalence implies C_1^s -equivalence. Hence $(C_1^s + C_n)$ -equivalence implies link-homotopy. By Corollary 4.5, L and L' are link-homotopic. $\square$

Remark 4.6. Let L be an n -component link in S^3 . Denote by $\mathcal{L}(L)$ the set of all n -component string links l such that $\text{cl}(l) = L$. Put $\mathcal{F}_2 \cup \cdots \cup \mathcal{F}_n = \{\pi_1, \dots, \pi_q\}$ so that any element in $\mathcal{F}_i$ appears before the elements in $\mathcal{F}_j$ ($2 \leq i < j \leq n$) and fix it. Then, by Remark 4.4, each l in $\mathcal{L}(L)$ is link-homotopic to $V_{\pi_1}^{x_{\pi_1}} * \cdots * V_{\pi_q}^{x_{\pi_q}}$ ($x_{\pi_k} = \mu_{\pi_k}(l) - \mu_{\pi_k}(\prod_{i=1}^{k-1} V_{\pi_i}^{x_{\pi_i}})$), which is the unique representation up to link-homotopy. We define a vector v_l as $v_l = (x_{\pi_1}, \dots, x_{\pi_q})$, and set $\mathcal{V}_L = \{v_l \mid l \in \mathcal{L}(L)\}$. By the uniqueness of the presentation for l , we have the following: *Two n -component links L and L' in S^3 are link-homotopic if and only if $\mathcal{V}_L \cap \mathcal{V}_{L'} \neq \emptyset$.*

5. SELF Δ -EQUIVALENCE OF BRUNNIAN LINKS

Let n and m be integers ($2 \leq n < m \leq 2n$). Given $k \in \{1, \dots, n\}$, consider a surjection τ from $\{1, \dots, m-2\}$ to $\{1, \dots, n\} \setminus \{k\}$. Let G_τ and $\overline{G}_\tau$ be simple C_{m-1}^a -trees illustrated in Figure 5.1, and set $V_\tau = (\mathbf{1}_n)_{G_\tau}$ and $V_\tau^{-1} = (\mathbf{1}_n)_{\overline{G}_\tau}$.




FIGURE 5.1. $\tau(i)$ and k mean $\tau(i)$ th and k th components of $\mathbf{1}_n$ respectively

Here, Figure 5.1 are the images of homeomorphisms from the neighborhoods of G_τ and $\overline{G}_\tau$ to the 3-ball. Although V_τ and V_τ^{-1} are not unique up to ambient isotopy, by Lemmas 2.1 and 2.3, they are unique up to C_m^a -equivalence. So, we may choose V_τ and V_τ^{-1} uniquely up to C_m^a -equivalence.

Set

$$\mu_\tau(L) = \mu_L(\tau(1) \dots \tau(m-2) \ k \ k).$$

Let $\mathcal{B}_m(k)$ be the set of all surjections τ from $\{1, \dots, m-2\}$ to $\{1, \dots, n\} \setminus \{k\}$ such that $|\tau^{-1}(i)| \leq 2$ ($i = 1, \dots, n$) and $|\tau^{-1}(j)| = 1$ (if $j > k$), and let ρ_m be a surjection

from $\{1, \dots, m-2\}$ to itself defined by $\rho_m(i) = m-1-i$. Note that the definition of $\mathcal{B}_m(k)$ implies that $\mathcal{B}_m(k) = \emptyset$ if $k < m-n$. So we may assume that $k \geq m-n$.

If $m \leq 2n-2$, then for any $\tau \in \mathcal{B}_m(k)$, $\{i \mid \tau(i) \neq \tau(m-1-i)\} \neq \emptyset$. We set

$$\mathcal{P}_m(k) = \left\{ \tau \in \mathcal{B}_m(k) \mid \begin{array}{l} \tau(p) < \tau(m-1-p) \\ \text{for } p = \min\{i \mid \tau(i) \neq \tau(m-1-i)\} \end{array} \right\}.$$

If $m = 2n-1$, then $k = n, n-1$ and there exists $\tau \in \mathcal{B}_{2n-1}(n) \cup \mathcal{B}_{2n-1}(n-1)$ such that $\tau(i) = \tau(2n-2-i)$ ($i = 1, \dots, n-2$) and $|\tau^{-1}(\tau(n-1))| = 1$. For $k = n, n-1$, set

$$\mathcal{R}_{2n-1}(k) = \left\{ \tau \in \mathcal{B}_{2n-1}(k) \mid \begin{array}{l} \tau(i) = \tau(2n-2-i) \ (i = 1, \dots, n-2), \\ |\tau^{-1}(\tau(n-1))| = 1 \end{array} \right\},$$

and set

$$\mathcal{P}_{2n-1}(k) = \left\{ \tau \in \mathcal{B}_{2n-1}(k) \setminus \mathcal{R}_{2n-1}(k) \mid \begin{array}{l} \tau(p) < \tau(2n-2-p) \\ \text{for } p = \min\{i \mid \tau(i) \neq \tau(2n-2-i)\} \end{array} \right\}.$$

Note that if $\tau \in \mathcal{R}_{2n-1}(n-1)$, then $\tau(n-1) = n$.

If $m = 2n$, then $k = n$ and there exists $\tau \in \mathcal{B}_{2n}(n)$ such that $\tau(i) = \tau(2n-1-i)$ ($i = 1, \dots, n-1$). Set

$$\mathcal{R}_{2n}(n) = \{\tau \in \mathcal{B}_{2n}(n) \mid \tau(i) = \tau(2n-1-i) \ (i = 1, \dots, n-1)\},$$

and set

$$\mathcal{P}_{2n}(n) = \left\{ \tau \in \mathcal{B}_{2n}(n) \setminus \mathcal{R}_{2n}(n) \mid \begin{array}{l} \tau(p) < \tau(2n-1-p) \\ \text{for } p = \min\{i \mid \tau(i) \neq \tau(2n-1-i)\} \end{array} \right\}.$$

We note that if $\tau \in \mathcal{R}_m(k)$ then $\tau\rho_m \in \mathcal{R}_m(k)$ (i.e., τ has ‘symmetry’), if $\tau \in \mathcal{P}_m(k)$ then $\tau\rho_m \notin \mathcal{P}_m(k)$, and

$$\mathcal{B}_m(k) = \mathcal{P}_m(k) \cup \mathcal{R}_m(k) \cup \{\tau\rho_m \mid \tau \in \mathcal{P}_m(k)\}.$$

For any $\varphi \in \mathcal{B}_m(k)$, V_φ is C_{m-1} -equivalent to $\mathbf{1}_n$. By Lemma 3.4, $\mu_{V_\varphi}(I) = 0$ for any I with $|I| \leq m-1$.

By the arguments similar to that in the proof of [16, Proposition 5.1], we have the following lemma.

Lemma 5.1. (1) If $\tau \in \mathcal{P}_m(k)$, then for an n -string link L ,

$$\text{cl}(L * V_{\tau\rho_m}) \stackrel{C_m^a}{\sim} \text{cl}(L * V_\tau) \text{ or } \text{cl}(L * V_\tau^{-1}).$$

Moreover the C_m^a -equivalence is realized by surgery along simple C_m^a -trees with $r_j \geq r_j(G_\tau)$ ($j = 1, \dots, n$).

(2) If $\varphi \in \mathcal{R}_{2n-1}(k)$, then for an n -string link L ,

$$\text{cl}(L * V_\varphi) \stackrel{C_{2n-1}^a}{\sim} \text{cl}(L * V_\varphi^{-1}).$$

Moreover the C_{2n-1}^a -equivalence is realized by surgery along simple C_m^a -trees with $r_j \geq r_j(G_\varphi)$ ($j = 1, \dots, n$).

Proof. (1) For the pairs of leaves grasping same components, by sliding the upper leaves on $\text{cl}(L)$ along the orientation, they come to below the others. Lemmas 2.1, 2.2 and 2.3 complete the proof. The same arguments give us a proof of (2). $\square$

Now we will calculate some Milnor invariants of string links V_φ for $\varphi \in \mathcal{R}_m(k) \cup \mathcal{P}_m(k)$.

Lemma 5.2. *For $\varphi \in \mathcal{P}_m(k)$ ($n+1 \leq m \leq 2n$, $m-n \leq k \leq n$) and $\tau \in \bigcup_l (\mathcal{R}_m(l) \cup \mathcal{P}_m(l))$,*

$$\mu_\tau(V_\varphi) = \begin{cases} 1 & \text{if } \varphi = \tau, \\ 0 & \text{if } \varphi \neq \tau. \end{cases}$$

Proof. We take the following 4 steps to proving this lemma.

Step 1: *Make a new link W_φ from V_φ by taking parallels of the components of V_φ so that Lemma 3.2 can be applied.*

Let $\tau \in \mathcal{R}_m(l) \cup \mathcal{P}_m(l)$ ($m-n \leq l \leq n$). Let $\mathbf{1}_m$ be the m -component trivial string link obtained from $\mathbf{1}_n$ by taking parallels of the components of $\mathbf{1}_n$ such that the i th component of $\mathbf{1}_m$ parallels to either

$$\begin{cases} \text{the } \tau(i)\text{th component of } \mathbf{1}_n \text{ if } i = 1, \dots, m-2, \text{ or} \\ \text{the } l\text{th component of } \mathbf{1}_n \text{ if } i = m-1, m, \end{cases}$$

and that $\mathbf{1}_m$ is contained in the tubular neighborhood $N(\mathbf{1}_n)$ of $\mathbf{1}_n$ with $G_\varphi \cap \mathbf{1}_m \subset \text{int}(G_\varphi \cap N(\mathbf{1}_n))$. Since a surgery along C_{m-1} -tree preserves framings, the above correspondance can be naturally extended so that the i th component of $(\mathbf{1}_m)_{G_\varphi}$ parallels to either

$$\begin{cases} \text{the } \tau(i)\text{th component of } (\mathbf{1}_n)_{G_\varphi} \text{ if } i = 1, \dots, m-2, \text{ or} \\ \text{the } l\text{th component of } (\mathbf{1}_n)_{G_\varphi} \text{ if } i = m-1, m. \end{cases}$$

Set $W_\varphi = (\mathbf{1}_m)_{G_\varphi}$. By Lemma 3.2, we have

$$\mu_{W_\varphi}(12\dots m) = \mu_{V_\varphi}(\tau(1)\dots\tau(m-2)ll)(= \mu_\tau(V_\varphi)).$$

Step 2: *By applying Lemma 2.5, deform W_φ up to the C_m -equivalence into*

$$(\mathbf{1}_m)_{G_{\varphi_1}} * (\mathbf{1}_m)_{G_{\varphi_2}} * \dots * (\mathbf{1}_m)_{G_{\varphi_s}}$$

so that each G_{φ_j} is a simple C_{m-1} -tree.

Set $V_j = (\mathbf{1}_m)_{G_{\varphi_j}}$ ($j = 1, \dots, s$).

Step 3: *By applying Lemmas 3.4 and 3.3, then we have*

$$\mu_{W_\varphi}(12\dots m) = \mu_{V_1}(12\dots m) + \dots + \mu_{V_s}(12\dots m).$$

Step 4: *If G_{φ_j} is a $C_{m-1}^{(2)}$ -tree, then by Proposition 2.9, V_j is link-homotopic to trivial, hence $\mu_{V_j}(12\dots m) = 0$. Otherwise, by using Lemma 4.1, calculate each $\mu_{V_j}(12\dots m)$.*

If $(|\varphi^{-1}(1)|, \dots, |\varphi^{-1}(n)|) \neq (|\tau^{-1}(1)|, \dots, |\tau^{-1}(n)|)$, then each G_{φ_j} is a $C_{m-1}^{(2)}$ -tree. This implies that $\mu_{W_\varphi}(12\dots m) = 0$.

Suppose $(|\varphi^{-1}(1)|, \dots, |\varphi^{-1}(n)|) = (|\tau^{-1}(1)|, \dots, |\tau^{-1}(n)|)$. Then each G_{φ_j} is a C_{m-1}^d -tree. Since $|\varphi^{-1}(k)| = 0$ and $|\tau^{-1}(i)| \geq 1$ ($i \neq l$), we have $k = l$, i.e., $\varphi \in \mathcal{P}_m(l)$.

If $\varphi \neq \tau$, then neither $(\varphi(1), \dots, \varphi(m-1))$ nor $(\varphi\rho_m(1), \dots, \varphi\rho_m(m-1))$ is equal to $(\tau(1), \dots, \tau(m-1))$. By Lemma 4.1, $\mu_{V_j}(12\dots m) = 0$ for any $j (= 1, 2, \dots, s)$.

If $\varphi = \tau$, then by Lemma 4.1 and by the fact that $\varphi\rho_m \neq \varphi$, there is a unique C_{m-1}^d -tree G_{φ_u} in $\{G_{\varphi_1}, \dots, G_{\varphi_s}\}$ such that

$$\mu_{V_j}(12\dots m) = \begin{cases} 1 & \text{if } j = u, \\ 0 & \text{if } j \neq u. \end{cases}$$

This completes the proof. $\square$

Remark 5.3. The calculation method used in the proof of Lemma 5.2 can be applied for another case. Let T be a linear, simple C_{m-1} -tree for $\mathbf{1}_n$ with the ends grasping k th component, and let $I = i_1 \dots i_{m-2} k k$ be a multi-index. Then, $\mu_{(\mathbf{1}_n)_T}(I)$ can be calculated as follows.

Step 1: Make a new link $W = (\mathbf{1}_m)_T$ from $(\mathbf{1}_n)_T$ by taking parallels of the components of $(\mathbf{1}_n)_T$ so that Lemma 3.2 can be applied.

Step 2: By applying Lemma 2.5, deform W up to the C_m -equivalence into $(\mathbf{1}_m)_{T_1} * (\mathbf{1}_m)_{T_2} * \dots * (\mathbf{1}_m)_{T_s}$ so that each T_j is a simple C_{m-1} -tree.

Step 3: By applying Lemmas 3.4 and 3.3, we have $\mu_W(12\dots m) = \mu_{(\mathbf{1}_m)_{T_1}}(12\dots m) + \dots + \mu_{(\mathbf{1}_m)_{T_s}}(12\dots m)$.

Step 4: If T_j is a $C_{m-1}^{(2)}$ -tree, then by Proposition 2.9, $\mu_{(\mathbf{1}_m)_{T_j}}(12\dots m) = 0$. Otherwise, by using Lemma 4.1, calculate each $\mu_{(\mathbf{1}_m)_{T_j}}(12\dots m)$.

Lemma 5.4. (1) For any $\varphi \in \mathcal{R}_{2n-1}(k)$ ($k = n, n-1$), the Milnor invariants of V_φ of length $\leq 2n-1$ vanish.

(2) For $\varphi \in \mathcal{R}_{2n-1}(n)$ and $\tau \in \mathcal{R}_{2n}(n) \cup \mathcal{P}_{2n}(n)$,

$$|\mu_\tau(V_\varphi)| = \begin{cases} 1 & \text{if } \tau \in \mathcal{R}_{2n}(n) \text{ and } \varphi(i) = \tau(i) \ (i = 1, \dots, n-1), \\ 0 & \text{otherwise.} \end{cases}$$

(3) For $\tau, \varphi \in \mathcal{R}_{2n}(n) \cup \mathcal{P}_{2n}(n)$,

$$\mu_\tau(V_\varphi) = \begin{cases} 1 & \text{if } \varphi = \tau \in \mathcal{P}_{2n}(n), \\ 2 & \text{if } \varphi = \tau \in \mathcal{R}_{2n}(n), \\ 0 & \text{if } \varphi \neq \tau. \end{cases}$$

Remark 5.5. (1) Note that, for any $\varphi \in \mathcal{R}_{2n-1}(n)$, there is a unique element $\tau \in \mathcal{R}_{2n}(n)$ such that $\varphi(i) = \tau(i)$ ($i = 1, \dots, n-1$), and that the correspondence induces a bijection from $\mathcal{R}_{2n-1}(n)$ to $\mathcal{R}_{2n}(n)$.

(2) For $\varphi \in \mathcal{R}_{2n-1}(n-1)$ and $\tau \in \mathcal{R}_{2n}(n) \cup \mathcal{P}_{2n}(n)$, while the Milnor invariants of V_φ of length $\leq 2n-1$ vanish, $\mu_\tau(V_\varphi)$ is not easily calculated. And we do not need the calculations to prove Theorem 1.3

Proof. As illustrated in Figure 5.2, the Whitehead link, which is a link $C(12, 12)$ defined in [1, subsection 7.11], is obtained from the 2-component trivial link by surgery along a simple C_2 -tree. We recall that, for a sequence $i_1 \dots i_k$,

(i) $C(i_1 i_2, i_1 i_2)$ is a Whitehead link; and

(ii) $C(i_1 \dots i_k i_{k+1}, i_1 \dots i_k i_{k+1}) = K_{i_1} \cup \dots \cup K_{i_{k+1}}$ is a link obtained from $C(i_1 \dots i_k, i_1 \dots i_k) = K_{i_1} \cup \dots \cup K_{i_{k-1}} \cup K'_{i_k}$ by replacing K'_{i_k} with Bing doubling $K_{i_k} \cup K_{i_{k+1}}$ of K'_{i_k} .

A 4-component link obtained from the 4-component trivial link by surgery along the 11 *basic claspers* illustrated in Figure 5.3 is ambient isotopic to a link illustrated in Figure 5.4, and to a link obtained from the trivial link by surgery along a clasper

with *boxes* as illustrated in Figure 5.5 (for the definitions a basic clasper and a box, see [10]). Since a link illustrated in Figure 5.4 is ambient isotopic to a link $C(1234, 1234)$, the link $C(1234, 1234)$ is C_7 -equivalent to $\text{cl}((\mathbf{1}_4)_G)$, where G is a simple C_6 -tree as illustrated in Figure 5.6. Moreover a band sum of $C(1234, 1234)$ and $\mathbf{1}_4$ as illustrated in Figure 5.7 is C_7 -equivalent to $(\mathbf{1}_4)_G$.

Similarly, we can see that, for $\varphi \in \mathcal{R}_{2n-1}(k)$ ($k = n, n-1$), a link $C(\alpha, \alpha)$ ($\alpha = \varphi(n-1)\varphi(n-2)\cdots\varphi(1)k$) is C_{2n-1} -equivalent to either $\text{cl}(V_\varphi)$ or $\text{cl}(V_\varphi^{-1})$. Then by Lemma 5.1 (2), $C(\alpha, \alpha)$ is C_{2n-1} -equivalent to $\text{cl}(V_\varphi)$.

In [1, subsection 7.11], it is shown that $\overline{\mu}_{C(\alpha, \alpha)}(I) = 0$ for any I with $|I| \leq 2n-1$. Hence, by Lemma 3.4, we have the conclusion (1).




FIGURE 5.2. (a): 2-component trivial link with a simple C_2 -clasper.
(b): 2-component trivial link with 3 basic claspers.




FIGURE 5.3. 4-component trivial link with 11 basic claspers. The numbers, 1,2,3, and 4, means the order of components.

Let $\tau \in \mathcal{R}_{2n}(n)$. Then it is not hard to see that $\text{cl}(V_\tau)$ is a link $L((\overline{\beta}, \beta))$ ($\beta = \tau(n-1)\tau(n-2)(\cdots(\tau(1)n)\cdots)$, $\overline{\beta} = ((\cdots(n\tau(1))\cdots)\tau(n-2))\tau(n-1)$) defined




FIGURE 5.4.




FIGURE 5.5.

in [1, subsection 7.4]. By combining [1, Proposition 6.5, Theorem 7.10 in p42, and Theorem 7.10 in p43], we have that if $\alpha = \beta$, i.e., $\varphi(i) = \tau(i)$ ($i = 1, \dots, n - 1$), then

$$|2\bar{\mu}_{C(\alpha, \alpha)}(I)| = |\bar{\mu}_{L((\bar{\beta}, \beta))}(I)|$$

for any I with $|I| = 2n$. Then, (2) follows from (3).

For $\tau, \varphi \in \mathcal{R}_{2n}(k) \cup \mathcal{P}_{2n}(n)$, by following the 4 steps in Remark 5.3, we have (3). This completes the proof. $\square$





FIGURE 5.6.

FIGURE 5.7. A band sum of $\mathbf{1}_4$ and $C(1234, 1234)$

Lemma 5.6. *Let L be an n -component string link and m ($n + 1 \leq m \leq 2n$) an integer. Let T be a simple C_{m-1}^a -tree for L . Suppose that $\mu_L(I) = 0$ for any I with $|I| \leq m - 1$ and $r(I) \leq 2$, and that T is not a simple $C_{m-1}^{(3)}$ -tree. Set $k = \max\{i \mid |T \cap (i\text{th component of } \mathbf{1}_n)| = 2\}$. Then*

(1) $\text{cl}(L_T)$ is C_m^a -equivalent to $\text{cl}(L * L')$, where $L' =$

$$\begin{cases} \prod_{\tau \in \mathcal{P}_m(k)} V_{\tau}^{\mu_{\tau}(L_T) - \mu_{\tau}(L)} & \text{if } n + 1 \leq m \leq 2n - 2, \\ \prod_{\tau \in \mathcal{P}_{2n-1}(k)} V_{\tau}^{\mu_{\tau}(L_T) - \mu_{\tau}(L)} * \prod_{\varphi \in \mathcal{R}_{2n-1}(k)} V_{\varphi}^{\varepsilon(\varphi)} & \text{if } m = 2n - 1, \\ \quad \quad \quad \text{(for some } \varepsilon(\varphi)\text{'s in } \{0, 1\}) \\ \prod_{\tau \in \mathcal{P}_{2n}(n)} V_{\tau}^{\mu_{\tau}(L_T) - \mu_{\tau}(L)} * \prod_{\varphi \in \mathcal{R}_{2n}(n)} V_{\varphi}^{(\mu_{\varphi}(L_T) - \mu_{\varphi}(L))/2} & \text{if } m = 2n, \end{cases}$$

and

(2) the C_m^a -equivalence is realized by surgery along simple C_m^a -trees with $r_k \geq 2$.

Note that if $m \geq n + 1$, then a simple C_{m-1}^a -tree is a simple $C_{m-1}^{(2)}$ -tree. If T is a simple $C_{m-1}^{(3)}$ -tree, then by Proposition 2.9, L_T and L are self Δ -equivalent.

Proof. By Lemma 2.1, L_T is C_m^a -equivalent to $L * (\mathbf{1}_n)_{T'}$, where T' is a simple C_{m-1}^a -tree and not a $C_{m-1}^{(3)}$ -tree with

$$\max\{i \mid |T' \cap (i\text{th component of } \mathbf{1}_n)| = 2\} = k.$$

By induction on the length of the path connecting two leaves which grasp k th component with applying Lemma 2.4, we have that $(\mathbf{1}_n)_{T'}$ is C_m^a -equivalent to a string

link which is obtained from $\mathbf{1}_n$ by surgery along simple linear C_{m-1}^a -trees whose ends grasp k th component of $\mathbf{1}_n$. By Lemmas 2.1, 2.2, 2.3, we have that

$$L_T \stackrel{C_m^a}{\sim} L * \prod_{\tau \in \mathcal{B}_m(k)} V_\tau^{x_\tau}.$$

Moreover, by Lemma 5.1, $\text{cl}(L * \prod_{\tau \in \mathcal{B}_m(k)} V_\tau^{x_\tau})$ is C_m^a -equivalent to $\text{cl}(L * L')$, where

$$L' = \begin{cases} \prod_{\tau \in \mathcal{P}_m(k)} V_\tau^{y_\tau} & \text{if } n+1 \leq m \leq 2n-2, \\ \prod_{\tau \in \mathcal{P}_m(k)} V_\tau^{y_\tau} * \prod_{\varphi \in \mathcal{R}_m(k)} V_\varphi^{\varepsilon(\varphi)}, \quad (\varepsilon(\varphi) \in \{0, 1\}) & \text{if } m = 2n-1, \\ \prod_{\tau \in \mathcal{P}_m(k)} V_\tau^{y_\tau} * \prod_{\varphi \in \mathcal{R}_m(k)} V_\varphi^{z_\varphi} & \text{if } m = 2n. \end{cases}$$

Note that the C_m^a -equivalences which are used in the above can be realized by surgery along simple C_m^a -trees with $r_k \geq 2$. By Lemmas 3.3, 5.2, and 5.4, for $\eta \in \mathcal{P}_m(k) \cup \mathcal{R}_m(k)$, we have that

$$\mu_\eta(L * L') = \begin{cases} \mu_\eta(L) + y_\eta & \text{if } \eta \in \mathcal{P}_m(k), \\ \mu_\eta(L) + 2z_\eta & \text{if } m = 2n \text{ and } \eta \in \mathcal{R}_m(k). \end{cases}$$

Since Milnor invariants of L with length $\leq m-1$ and $r \leq 2$ vanish, by Lemma 3.4, those of L_T also vanish. By combining this, the fact that $\text{cl}(L_T)$ and $\text{cl}(L * L')$ are C_m^a -equivalence, and Lemma 3.4, we have that

$$\mu_\eta(L_T) = \bar{\mu}_\eta(\text{cl}(L_T)) = \bar{\mu}_\eta(\text{cl}(L * L')) = \mu_\eta(L * L').$$

This completes the proof. $\square$

The following is the main result in this section.

Theorem 5.7. *Let L be an n -component Brunnian link. If $\bar{\mu}_L(I) = 0$ for any I with $|I| \leq 2n-1$ and $r(I) \leq 2$, then L is self Δ -equivalent to the closure of $L' * L''$, where*

$$L' = \prod_{\varphi \in \mathcal{R}_{2n-1}(n)} V_\varphi^{\varepsilon(\varphi)}, \quad L'' = \prod_{\tau \in \mathcal{R}_{2n}(n)} V_\tau^{(\bar{\mu}_\tau(L) - \bar{\mu}_\tau(L'))/2} * \prod_{\eta \in \mathcal{P}_{2n}(n)} V_\eta^{\bar{\mu}_\eta(L)},$$

and

$$\varepsilon(\varphi) = \begin{cases} 1 & \text{if } \bar{\mu}_\tau(L) \text{ is odd for } \tau \in \mathcal{R}_{2n}(n) \text{ with } \tau(i) = \varphi(i) \ (i = 1, \dots, n-1) \\ 0 & \text{if } \bar{\mu}_\tau(L) \text{ is even for } \tau \in \mathcal{R}_{2n}(n) \text{ with } \tau(i) = \varphi(i) \ (i = 1, \dots, n-1). \end{cases}$$

Note that, in the theorem above, $L' * L''$ is determined by Milnor invariants of L with length $2n$ and $r = 2$.

Proof. By Proposition 2.6, L is obtained from the n -component trivial link O by surgery along simple C_{n-1}^a -trees $T_1, \dots, T_l$. Hence we have

$$L = \text{cl}((\mathbf{1}_n)_{T_1 \cup T_2 \cup \dots \cup T_l}).$$

By Lemmas 4.2, 2.1, 2.2 and 2.3, we have that

$$L \stackrel{C_n^a}{\sim} \text{cl}\left(\prod_{\pi \in \mathcal{F}_n} V_\pi^{\bar{\mu}_\pi(L)}\right).$$

Since $\bar{\mu}_\pi(L) = 0$ for any $\pi \in \mathcal{F}_n$, L is C_n^a -equivalent to O , i.e., L is obtained from O by surgery along simple C_n^a -trees. By Lemmas 5.6 (1), 2.1, 2.2 and 2.3, we have

that

$$L \stackrel{C_{n+1}^a}{\sim} \text{cl} \left(\prod_{1 \leq k \leq n} \left(\prod_{\tau \in \mathcal{P}_{n+1}(k)} V_{\tau}^{\bar{\mu}_{\tau}(L)} \right) \right).$$

Since $\bar{\mu}_{\tau}(L) = 0$ for any $\tau \in \mathcal{P}_{n+1}(k)$ ($k = 1, \dots, n$), L is C_{n+1}^a -equivalent to O . Note that a simple C_m^a -tree ($m \geq n+1$) for an n -component link is a simple $C_m^{(2)}$ -tree and might be $C_m^{(3)}$ -tree. By Lemmas 5.6 (1), 2.1, 2.2 and 2.3, we have that

$$L \stackrel{C_2^s + C_{n+2}^a}{\sim} \text{cl} \left(\prod_{2 \leq k \leq n} \left(\prod_{\tau \in \mathcal{P}_{n+2}(k)} V_{\tau}^{\bar{\mu}_{\tau}(L)} \right) \right).$$

Since $\bar{\mu}_{\tau}(L) = 0$ for any $\tau \in \mathcal{P}_{n+2}(k)$ ($k = 2, \dots, n$), by repeating this step, then by Lemma 5.1 (2), we have that

$$L \stackrel{C_2^s + C_{2n-1}^a}{\sim} \prod_{\varphi \in \mathcal{R}_{2n-1}(n)} V_{\varphi}^{\varepsilon(\varphi)} * \prod_{\phi \in \mathcal{R}_{2n-1}(n-1)} V_{\phi}^{\varepsilon(\phi)}$$

for some $\varepsilon(\varphi)$'s and $\varepsilon(\phi)$'s in $\{0, 1\}$.

In the proof of Lemma 5.4 (1), we showed that, for $\phi \in \mathcal{R}_{2n-1}(n-1)$,

$$\text{cl}(V_{\phi}) \stackrel{C_{2n-1}}{\sim} C(\alpha, \alpha),$$

where $\alpha = n\phi_j(n-2) \cdots \phi_j(1)(n-1)$. Since the Whitehead link $C(12, 12)$ is deformed into a trivial link by a single self crossing change in the first component, $C(\alpha, \alpha)$ is also deformed into a trivial link by a single self crossing change in the n th component. So $C(\alpha, \alpha)$ is obtained from a trivial link by surgery along a simple C_1^s -tree T with $r_n(T) = 2$. By Lemma 2.7, $C(\alpha, \alpha)$ is obtained from a trivial link by surgery along simple C_n^a -trees with $r_n = 2$. Since the Milnor invariants of $C(\alpha, \alpha)$ with length $\leq 2n-1$ vanish, by Lemmas 5.6, 2.1, 2.2, 2.3 and 5.1 (2), we have that

$$C(\alpha, \alpha) \stackrel{C_2^s + C_{2n-1}^a}{\sim} \text{cl} \left(\prod_{\varphi \in \mathcal{R}_{2n-1}(n)} V_{\varphi}^{\varepsilon'(\varphi)} \right)$$

for some $\varepsilon'(\varphi)$'s in $\{0, 1\}$.

Since

$$\prod_{\varphi \in \mathcal{R}_{2n-1}(n)} V_{\varphi}^{\varepsilon(\varphi)} * \prod_{\phi \in \mathcal{R}_{2n-1}(n-1)} V_{\phi}^{\varepsilon(\phi)}$$

is obtained from $\mathbf{1}_n$ by surgery along simple C_{2n-2}^a -trees, by Lemmas 2.1, 2.2 and 5.1 (2), L is $(C_2^s + C_{2n-1})$ -equivalent to the closure of

$$L' = \prod_{\varphi \in \mathcal{R}_{2n-1}(n)} V_{\varphi}^{\varepsilon''(\varphi)},$$

where $\varepsilon''(\varphi)$'s are integers in $\{0, 1\}$.

Note that a simple C_{2n-1} -tree for an n -component link is either C_{2n-1}^a -tree or $C_{2n-1}^{(3)}$ -tree, hence by Proposition 2.9, C_{2n-1} -equivalence implies $(C_2^s + C_{2n-1}^a)$ -equivalence. So L is self Δ -equivalent to a link obtained from $\text{cl}(L')$ by surgery along simple C_{2n-1}^a -trees. By Lemmas 5.6 (1), 2.1, 2.2 and 2.3, L is $(C_2^s + C_{2n}^a)$ -equivalent to $\text{cl}(L' * L'')$, where

$$L'' = \prod_{\tau \in \mathcal{R}_{2n}(n)} V_{\tau}^{(\mu_{\tau}(L) - \mu_{\tau}(L'))/2} * \prod_{\eta \in \mathcal{P}_{2n}(n)} V_{\eta}^{\mu_{\eta}(L)}.$$

Since a C_{2n} -tree is a $C_{2n}^{(3)}$ -tree, by Proposition 2.9, L is self Δ -equivalent to the closure of $L' * L''$. By Lemmas 3.3, 3.4 and 5.4, we have that for any $\tau \in \mathcal{R}_{2n-1}(n)$,

$$\overline{\mu}_\tau(L) \equiv \varepsilon''(\varphi) \pmod{2},$$

where $\varphi(i) = \tau(i)$ ($i = 1, \dots, n-1$). This completes the proof. $\square$

By Theorem 5.7, we have the following two corollaries. Corollaries 5.8 and 5.9 are special cases of Theorem 1.3 and Corollary 1.5, respectively. Since these corollaries are needed to show Theorem 1.3, we give the statements.

Corollary 5.8. *Let L and L' be n -component Brunnian links. Suppose that $\overline{\mu}_L(I) = \overline{\mu}_{L'}(I) = 0$ for any I with $|I| \leq 2n-1$ and $r(I) \leq 2$. Then L and L' are self Δ -equivalent if and only if $\overline{\mu}_L(J) = \overline{\mu}_{L'}(J)$ for any J with $|J| = 2n$ and $r(J) = 2$.*

Corollary 5.9. *A Brunnian link L is self Δ -equivalent to a trivial link if and only if $\overline{\mu}_L(I) = 0$ for any I with $r(I) \leq 2$.*

6. LINKS WITH MILNOR INVARIANTS VANISH

Before proving Theorem 1.3, we need some preparations.

Let $L = K_1 \cup \dots \cup K_n$ be an n -component link and b a band attaching a single component K_i with orientation coherent, i.e., $b \cap L = K_i \cap b \subset \partial b$ consists of two arcs whose orientations from K_i are opposite to those from ∂b . Then the $(n+1)$ -component link $L' = (L \cup \partial b) \setminus \text{int}(b \cap K_i)$ is called a link obtained from L by *fission* (along a band b), and conversely L is called a link obtained from L' by *fusion* [13]. Let $L' = K_{11} \cup \dots \cup K_{1l_1} \cup \dots \cup K_{n1} \cup \dots \cup K_{nl_n}$ be a link obtained from an n -component link $K_1 \cup \dots \cup K_n$ by a finite sequence of fission, where $K_{i1} \cup \dots \cup K_{il_i}$ is obtained from K_i ($i = 1, \dots, n$). We assign a color $c(K_{ij})$ to K_{ij} as $c(K_{ij}) = i$. In this section, for a C_k -tree T , we call T a C_k^s -tree (resp. C_k^d -tree) if $|\{c(K_{ij}) \mid T \cap K_{ij} \neq \emptyset\}| = 1$ (resp. $= k+1$). A C_k^s -move (resp. C_k^d -move) is a local move defined by surgery along simple C_k^s -tree (resp. C_k^d -tree).

Lemma 6.1. *If an n -component link is deformed into a trivial link by a finite sequence of fission, C_2^s -moves and C_{n-1}^d -moves, then L is self Δ -equivalent to a Brunnian link.*

Proof. Note that L is obtained from a trivial link by a finite sequence of fusion, C_2^s -moves and C_{n-1}^d -moves. By the arguments similar to that in the proof of [10, Proposition 3.22], we may assume that the bands of fusion, C_2^s -trees and C_{n-1}^d -trees are mutually disjoint. So there exist an n -component ribbon link L_0 and a disjoint union $F \cup F'$ of simple C_2^s -trees and C_{n-1}^d -trees such that $L = L_0 \cup F \cup F'$, where F (resp. F') is a disjoint union of C_2^s -trees (resp. C_{n-1}^d -trees). Since ribbon links are self Δ -equivalent to a trivial link [24], L_0 is self Δ -equivalent to the n -component trivial link O . Hence

$$L \stackrel{C_2^s + C_{n-1}^d}{\sim} O.$$

This implies that L is self Δ -equivalent to a link obtained from O by surgery along simple C_{n-1}^d -trees. Since a C_{n-1}^d -tree is a C_{n-1}^a -tree, by Proposition 2.6, we have the conclusion. $\square$

Theorem 6.2. *Let L be an n -component link such that $\bar{\mu}_L(I) = 0$ for any I with $|I| \leq 2n - 2$ and $r(I) \leq 2$. Then L is self Δ -equivalent to a Brunnian link.*

Proof. By Lemma 6.1, it is enough to show that L is deformed into a trivial link by a finite sequence of fission, C_2^s -moves and C_{n-1}^d -moves.

Since any knot is Δ -equivalent to be trivial [20], we may assume that every component of L is trivial.

Suppose that any k -component sublink of L is Brunnian ($2 \leq k \leq n - 1$). Since surgery along a simple C_{k-1}^d -tree with index $\{i_1, \dots, i_k\}$ does not change the link type of $K_{j_1} \cup \dots \cup K_{j_k}$ for $\{j_1, \dots, j_k\} \neq \{i_1, \dots, i_k\}$, by Proposition 2.6, L is C_{k-1}^d -equivalence to a link L' whose the k -component sublinks are trivial. Let L_0 be an n -component string link with $\text{cl}(L_0) = L'$, and set $\mathbf{1}_n = \gamma_1 \cup \dots \cup \gamma_n$. Let $\{S_1, \dots, S_m\}$ ($m = \binom{n}{k}$) be the set of subsets of $\{1, \dots, n\}$ with $|S_i| = k$ ($i = 1, \dots, m$). By Lemmas 2.1 and 2.2,

$$L \stackrel{C_k}{\sim} \text{cl}(L_0 * L_1 * \dots * L_m),$$

where L_i is a string link which is splitable into the unions L_{i1} of the j th ($j \in S_i$) components and the other components L_{i0} such that L_{i1} is obtained from $\bigcup_{j \in S_i} \gamma_j$ by surgery along simple C_{k-1}^d -trees with indices S_i and $L_{i0} = \mathbf{1}_n - \bigcup_{j \in S_i} \gamma_j$, and the C_k -equivalence is realized by surgery along simple C_k -trees with $|\text{index}| \geq k$. This implies that L is C_k^d -equivalent to a link obtained from $\text{cl}(L_0 * L_1 * \dots * L_m)$ by surgery along simple C_k -trees with $|\text{index}| = k$.

By Lemmas 2.1 and 2.2,

$$L \stackrel{C_k^d + C_{k+1}}{\sim} \text{cl}(L_0 * L_1^1 * \dots * L_m^1),$$

where L_i^1 is a string link which is splitable into the unions L_{i1}^1 of the j th ($j \in S_i$) components and L_{i0} such that L_{i1}^1 is obtained from $\bigcup_{j \in S_i} \gamma_j$ by surgery along simple C_{k-1}^d , C_k -trees with indices S_i ($i = 1, \dots, m$), and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with $|\text{index}| \geq k$. By Proposition 2.8, surgery along a simple C_{k+1} -tree with $|\text{index}| \geq k + 1$ is realized by C_k^d -equivalence. Therefore, L is C_k^d -equivalent to a link obtained from $\text{cl}(L_0 * L_1^1 * \dots * L_m^1)$ by surgery along simple C_{k+1} -trees with $|\text{index}| = k$.

By Lemmas 2.1 and 2.2,

$$L \stackrel{C_k^d + C_{k+2}}{\sim} \text{cl}(L_0 * L_1^2 * \dots * L_m^2),$$

where L_i^2 is a string link which is splitable into the unions L_{i1}^2 of the j th ($j \in S_i$) components and L_{i0} such that L_{i1}^2 is obtained from $\bigcup_{j \in S_i} \gamma_j$ by surgery along simple C_{k-1}^d , C_k , C_{k+1} -trees with indices S_i ($i = 1, \dots, m$), and the C_{k+2} -equivalence is realized by surgery along simple C_{k+2} -trees with $|\text{index}| \geq k$.

By repeating this procedure, we have that

$$L \stackrel{C_k^d + C_{2k}}{\sim} \text{cl}(L_0 * L_1^k * \dots * L_m^k),$$

where L_i^k is a string link which is splitable into the unions L_{i1}^k of the j th ($j \in S_i$) components and L_{i0} such that L_{i1}^k is obtained from $\bigcup_{j \in S_i} \gamma_j$ by surgery along simple

$C_{k-1}^d, C_k, \dots, C_{2k-1}$ -trees with indices S_i ($i = 1, \dots, m$). Note that a simple C_{2k} -tree is either a C_{2k} -tree with index $\geq k + 1$ or a $C_{2k}^{(3)}$ -tree. By Lemma 2.9,

$$L \stackrel{C_k^d + C_2^s}{\sim} \text{cl}(L_0 * L_1^k * \dots * L_m^k).$$

So L is self Δ -equivalent to a link obtained from $\text{cl}(L_0 * L_1^k * \dots * L_m^k)$ by surgery along simple C_k^d -trees. By Lemmas 2.1 and 2.2,

$$L \stackrel{C_2^s + C_{k+1}}{\sim} \text{cl}(M_1 * L_0 * L_1^k * \dots * L_m^k),$$

where M_1 is a string link obtained from $\mathbf{1}_n$ by surgery along simple C_k^d -trees, and the C_{k+1} -equivalence is realized by surgery along simple C_{k+1} -trees with $|\text{index}| \geq k + 1$.

By repeating this step, we have that

$$L \stackrel{C_2^s + C_{2n}}{\sim} \text{cl}(M_{2n-k} * L_0 * L_1^k * \dots * L_m^k),$$

where M_{2n-k} is a string link obtained from $\mathbf{1}_n$ by surgery along simple $C_k^d, C_{k+1}, \dots, C_{2n-1}$ -trees with $|\text{index}| \geq k + 1$. Since a simple C_{2n} -tree is a $C_{2n}^{(3)}$ -tree, by Lemma 2.9,

$$L \stackrel{C_2^s}{\sim} \text{cl}(M_{2n-k} * L_0 * L_1^k * \dots * L_m^k).$$

Note that $\text{cl}(M_{2n-k} * L_0 * L_1^k * \dots * L_m^k)$ is deformed into a split sum of $\text{cl}(M_{2n-k} * L_0), \text{cl}(L_{11}^k), \dots, \text{cl}(L_{m1}^k)$ by a finite sequence of fission. Since for any index J in S_i with $|J| \leq 2k (\leq 2n - 2)$ and $r(J) \leq 2$, $\mu_{L_{i1}^k}(J) = \bar{\mu}_L(J) = 0$, by Corollary 5.9, $\text{cl}(L_{i1}^k)$ is self Δ -equivalent to a trivial link.

Hence L is deformed into a split sum of $\text{cl}(M_{2n-k} * L_0)$ and a trivial link by a finite sequence of fission and C_2^s -moves. Note that any $(k + 1)$ -component sublink of $\text{cl}(M_{2n-k} * L_0)$ is Brunnian.

By the induction, we have that L is deformed into a split sum of an n -component Brunnian link B and a trivial link by a finite sequence of fission and self Δ -moves. By Proposition 2.6, B is C_{n-1}^d -equivalent to a trivial link. This completes the proof. $\square$

By combining Corollarie 5.8 and Theorem 6.2, we can prove Theorem 1.3.

Proof of Theorem 1.3. Let L be an n -component link with $\bar{\mu}_L(I) = 0$ for any I with $|I| \leq 2n - 1$ and $r(I) \leq 2$. By Theorem 6.2, L is self Δ -equivalent to a Brunnian link B . Since $\bar{\mu}_B(I) = \bar{\mu}_L(I) = 0$ for any I with $|I| \leq 2n - 1$ and $r(I) \leq 2$, by Corollary 5.8, B is determined by Milnor invariants with length $2n$ and $r = 2$. This completes the proof. $\square$

The following theorem characterizes n -component links whose Milnor invariants of length $\leq 2n - 1$ and $r \leq 2$ vanish.

Theorem 6.3. *For an n -component link L , $\bar{\mu}_L(I) = 0$ for any I with $|I| \leq 2n - 1$ and $r(I) \leq 2$ if and only if, for each $i \in \{1, \dots, n\}$, there is a Brunnian link L_i such that L_i is self Δ -equivalent to L and the i th component K of L_i is null-homotopic in $S^3 \setminus (L_i - K)$.*

Proof. For the ‘only if’ part, it is enough to consider the case when $i = n$. By Theorem 6.2, L is self Δ -equivalent to a Brunnian link. By Theorem 5.7, the Brunnian link is self Δ -equivalent to the closure L_n of a product of some V_φ ’s ($\varphi \in \mathcal{R}_{2n-1}(n) \cup \mathcal{R}_{2n}(n) \cup \mathcal{P}_{2n}(n)$). Note that, for $\varphi \in \mathcal{R}_{2n-1}(n)$ (resp. $\varphi \in \mathcal{R}_{2n}(n) \cup \mathcal{P}_{2n}(n)$)

V_φ is $C_{2n-1}^{(2)}$ -equivalent (resp. $C_{2n}^{(2)}$ -equivalent) to $\mathbf{1}_n$ and the $C_{2n-1}^{(2)}$ -equivalence (resp. $C_{2n}^{(2)}$ -equivalence) is realized by surgery along simple $C_{2n-1}^{(2)}$ -trees (resp. $C_{2n}^{(2)}$ -trees) with $r_n = 2$. By Proposition 2.9, L_n is self C_1 -equivalent to a trivial link and the self C_1 -equivalence is realized by surgery along simple C_1^s -trees with $r_n = 2$. Hence the n th component K of L_n is null-homotopic in $S^3 \setminus (L_n - K)$.

Now we will show the ‘if’ part. Let I be an index with $|I| \leq 2n - 1$ and $r(I) \leq 2$. Since L is self Δ -equivalent to a Brunnian link, if I does not contain an integer in $\{1, \dots, n\}$, then $\bar{\mu}_L(I) = 0$. So we may suppose that I contains any integer in $\{1, \dots, n\}$. The condition $|I| \leq 2n - 1$ implies that there is an integer i such that i appears in I once. Let L_i be a Brunnian link such that L_i is self Δ -equivalent to L and the i th component K of L_i is null-homotopic in $S^3 \setminus (L_i - K)$. This implies that $\bar{\mu}_{L_i}(Ji) = 0$ for any index J in $\{1, \dots, n\} \setminus \{i\}$. Since $\bar{\mu}$ has ‘cyclic symmetry’ ([18, Theorem 8]), $\bar{\mu}_{L_i}(I) = 0$. This completes the proof. $\square$

Example 6.4. Let V be a string link illustrated in Figure 6.1 and L be the closure of V . By Proposition 2.9, for i ($i = 2, 3$), V is self C_1 -equivalent to $\mathbf{1}_3$ and the self C_1 -equivalence is realized by surgery along C_1^s -trees with indices $\{i\}$. Hence the i th component K_i of L is null-homotopic in $S^3 \setminus (L - K_i)$ ($i = 2, 3$). Suppose that the 1st component K_1 is null-homotopic in $S^3 \setminus (L - K_1)$. Then, by Theorem 6.3, $\bar{\mu}_L(I) = 0$ for any I with $|I| \leq 5$ and $r(I) \leq 2$. By Lemma 5.4 (3), $\mu_V(12233) = 1$. Hence $\bar{\mu}_L(12233) = 1$. This is a contradiction.




FIGURE 6.1.

REFERENCES

- [1] T. Cochran, *Derivatives of links: Milnor’s concordance invariants and Massey’s products*, Mem. Amer. Math. Soc. **84** (1990), no. 427, x+73 pp.
- [2] J. Conant, P. Teichner, *Grope cobordism of classical knots*, Topology **43** (2004), 119–156.
- [3] T. Fleming and A. Yasuhara, *Milnor’s isotopy invariants and generalized link homotopy*, preprint math.GT/0511477.
- [4] T. Fleming and A. Yasuhara, *Milnor numbers and the self delta classification of 2-string links*, to appear in the proceedings of Intelligence of Low Dimensional Topology 2006, Knots and Everything Series, World Scientific.
- [5] C. H. Giffen, *Link concordance implies link homotopy*, Math. Scand. **45** (1979), 243–254.
- [6] D. L. Goldsmith, *Concordance implies homotopy for classical links in M^3* , Comment. Math. Helv. **54** (1979), 347–355.
- [7] M.N. Gusarov, *Variations of knotted graphs. The geometric technique of n -equivalence*. (Russian), Algebra i Analiz **12** (2000), no. 4, 79–125; translation in St. Petersburg Math. J. **12** (2001), no. 4, 569–604.

- [8] N. Habegger and X.S. Lin, *The classification of links up to link-homotopy*, J. Amer. Math. Soc. **3** (1990), 389–419.
- [9] K. Habiro, Master's thesis, University of Tokyo, 1994.
- [10] K. Habiro, *Claspers and finite type invariants of links*, Geom. Topol. **4** (2000), 1–83.
- [11] K. Habiro, *Brunnian links, claspers, and Goussarov-Vassiliev finite type invariants*, to appear in Math. Proc. Cambridge Philos. Soc.
- [12] K. Habiro and J.-B. Meilhan *Finite type invariants and Milnor invariants for Brunnian links*, preprint math.GT/0510534.
- [13] A. Kawauchi, T. Shibuya and S. Suzuki, Descriptions on surfaces in four space, I. Normal forms, Math. Sem. Notes, Kobe Univ. **10** (1982), 75–125.
- [14] J.-B. Meilhan, *Invariants de type fini des cylindres d'homologie et des string links*, Thèse de Doctorat (2003), Université de Nantes.
- [15] J.-B. Meilhan, *On surgery along Brunnian links in 3-manifolds*, Alg. Geom. Topol. **6** (2006), 2417–2453.
- [16] J.-B. Meilhan and A. Yasuhara, *On C_n -moves for links*, preprint math.GT/0607116
- [17] J. Milnor, *Link groups*, Ann. of Math. (2) **59** (1954), 177–195.
- [18] J. Milnor, *Isotopy of links*, Algebraic geometry and topology, A symposium in honor of S. Lefschetz, pp. 280–306, Princeton University Press, Princeton, N. J., 1957.
- [19] H.A. Miyazawa and A. Yasuhara, *Classification of n -component Brunnian links up to C_n -move*, Topology Appl. **153** (2006), 1643–1650.
- [20] H. Murakami and Y. Nakanishi, *On a certain move generating link-homology*, Math. Ann. **283** (1989), 75–89.
- [21] Y. Nakanishi and Y. Ohyama, *Delta link homotopy for two component links. III*, J. Math. Soc. Japan **55** (2003), 641–654.
- [22] Y. Nakanishi and T. Shibuya, *Link homotopy and quasi self delta-equivalence for links*, J. Knot Theory Ramifications **9** (2000), 683–691.
- [23] Y. Nakanishi, T. Shibuya and A. Yasuhara, *Self delta-equivalence of cobordant links*, Proc. Amer. Math. Soc. **134** (2006), 2465–2472.
- [24] T. Shibuya, *Self Δ -equivalence of ribbon links*, Osaka J. Math. **33** (1996), 751–760.
- [25] T. Shibuya and A. Yasuhara, *Self C_k -move, quasi self C_k -move and the Conway potential function for links*, J. Knot Theory Ramifications **13** (2004), 877–893.
- [26] T. Shibuya and A. Yasuhara, *Boundary links are self delta-equivalent to trivial links*, to appear in Math. Proc. Cambridge Philos. Soc.
- [27] N. Smythe, *Boundary links*, Topology Seminar (Univ. Wisconsin, Madison, Wis., 1965), pp. 69–72, Ann. of Math. Studies, No. 60, Princeton Univ.
- [28] K. Taniyama and A. Yasuhara, *Band description of knots and Vassiliev invariants*, Math. Proc. Cambridge Philos. Soc., **133** (2002), 325–343.

TOKYO GAKUGEI UNIVERSITY, DEPARTMENT OF MATHEMATICS, KOGANEISHI, TOKYO 184-8501, JAPAN

E-mail address: yasuhara@u-gakugei.ac.jp